



Contents lists available at ScienceDirect

Neuroscience Research

journal homepage: www.elsevier.com/locate/neures



Review article

New frontiers in probing the dynamics of purinergic transmitters *in vivo*

Zhaofa Wu^{a,b,*}, Yulong Li^{a,b,c,d,*}

^a State Key Laboratory of Membrane Biology, Peking University School of Life Sciences, Beijing, 100871, China

^b PKU-IDG/McGovern Institute for Brain Research, Beijing, 100871, China

^c Peking-Tsinghua Center for Life Sciences, Beijing, 100871, China

^d Chinese Institute for Brain Research, Beijing, 100871, China

ARTICLE INFO

Article history:

Received 13 December 2019

Received in revised form 3 January 2020

Accepted 5 January 2020

Available online xxx

Keywords:

Purinergic transmitters

ATP

Adenosine

Sensors

Imaging

ABSTRACT

Purinergic transmitters such as adenosine, ADP, ATP, UTP, and UDP-glucose play important roles in a wide range of physiological processes, including the sleep-wake cycle, learning and memory, cardiovascular function, and the immune response. Moreover, impaired purinergic signaling has been implicated in various pathological conditions such as pain, migraine, epilepsy, and drug addiction. Examining the function of purinergic transmission in both health and disease requires direct, sensitive, non-invasive tools for monitoring structurally similar purinergic transmitters; ideally, these tools should have high spatial and temporal resolution in *in vivo* applications. Here, we review the recent progress with respect to the development and application of new methods for detecting purinergic transmitters, focusing on optical tools; in addition, we provide discussion regarding future perspectives.

© 2020 Elsevier B.V. and Japan Neuroscience Society. All rights reserved.

Contents

1. Introduction.....	00
2. Non-optical methods for probing purinergic transmitters.....	00
2.1. Electrophysiology.....	00
2.2. Microdialysis.....	00
2.3. Amperometric biosensors.....	00
2.4. Fast-scan cyclic voltammetry.....	00
3. Optical methods.....	00
3.1. Optical sensors for detecting intracellular purine metabolites.....	00
3.2. Optical sensors for detecting extracellular purinergic transmitters.....	00
3.2.1. Bioluminescence-based sensors.....	00
3.2.2. Fluorescence-based sensors.....	00
4. Future perspectives.....	00
Author contributions.....	00
Acknowledgments and Declaration of Competing Interest.....	00
References.....	00

* Corresponding author at: State Key Laboratory of Membrane Biology, Peking University School of Life Sciences, Beijing, 100871, China.
E-mail addresses: wuzhaofa@pku.edu.cn (Z. Wu), yulongli@pku.edu.cn (Y. Li).

1. Introduction

Purines and pyrimidines are essential components in living cells, where they mediate energy conversion and form the basis for synthesizing nucleic acids. In addition to these intracellular functions, some purines and pyrimidines such as ATP, adenosine, UTP, and UDP-glucose—known collectively as purinergic transmitters—also play a key role in extracellular signaling via purinergic receptors (Ralevic and Burnstock, 1998). Among these transmitters, adenosine (Ado) and ATP are arguably two of the most thoroughly studied.

As early as 1929, Alan Drury and Albert Szent-Györgyi found that an intravenous injection of Ado induced sleep in animals (Drury and Szent-Györgyi, 1929), providing the first physiological description for this nucleotide. Subsequently, a role for ATP in chemical transmission was identified in sensory nerve endings (Holton and Holton, 1954; Holton, 1959), later giving rise to the concept of purinergic transmission after ATP was identified as the transmitter involved in non-adrenergic, non-cholinergic inhibitory nerves in guinea pig intestinal smooth muscle (Burnstock, 1972).

After decades of research, it is now widely accepted that ATP can function as a neurotransmitter (Pankratov et al., 2006), gliotransmitter (Maienschein et al., 1999; Zhang et al., 2003), or co-transmitter (Burnstock, 2007; Jo and Schlichter, 1999) upon release from secretory vesicles such as synaptic vesicles, glial vesicles, and lysosomes (Zhang et al., 2007) via exocytosis (MacDonald et al., 2006). In addition, ATP can be released from cells via connexin/pannexin hemichannels (Cotrina et al., 1998), anion channels (Sabiroy and Okada, 2005) and voltage-gated ion channels (Ma et al., 2018; Taruno et al., 2013). Extracellular Ado is believed to be generated by the breakdown of extracellular nucleotides, particularly ATP and ADP (Chen et al., 2013), via ecto-nucleotidases (Zimmermann et al., 2012), or by the direct efflux of Ado through the so-called equilibrative nucleoside transporters (ENTs) (Lovatt et al., 2012); whether Ado functions as a neurotransmitter released via the exocytosis of synaptic vesicle is currently under debate (Corti et al., 2013; Dale and Frenguelli, 2009). In the extracellular space, Ado and ATP perform distinct functions by activating P1 receptors and P2 receptors, respectively. The P1 receptor family consists of the A₁, A_{2A}, A_{2B}, and A₃ G protein-coupled receptors (GPCRs) (Chen et al., 2013), whereas P2 receptors include the ionotropic P2X receptor (Khakh and North, 2006) and the G protein-coupled metabotropic P2Y receptor. Both P1 and P2 receptors are widely expressed and have been implicated in a variety of physiological processes (Burnstock, 2008; Chen et al., 2013), including controlling the sleep-wake cycle (Huang et al., 2005; Porkka-Heiskanen et al., 1997), learning and memory (Dias et al., 2013), cardiovascular function (Shryock and Belardinelli, 1997), and the immune response (Eltzschig et al., 2012), as well as pathological processes, including pain (Hasko et al., 2008), epilepsy (Boison, 2012; During and Spencer, 1992; Engel et al., 2016), ischemia–reperfusion (Eltzschig and Eckle, 2011), drug addiction (Dunwiddie, 1999), and neurodegenerative disorders (Schwarzschild et al., 2006).

In addition to the broad effects that Ado and ATP have on various systems, functions of other purinergic transmitters are also identified. For example, ADP plays an important role in platelet aggregation by acting at P2Y₁ receptor (Jin et al., 1998); UTP has been found to play a role in nociception in the central (Li et al., 2014; Ren et al., 2016) and enteric (Hockley et al., 2016) nervous systems and is an agonist of P2Y receptors; UDP-glucose may induce inflammatory effects by acting at the P2Y₁₄ receptor (Lazarowski and Harden, 2015).

Despite the significant contributions to our understanding of purinergic signaling by many researchers over the past several decades, many key questions remain, including which purinergic transmitters are released—and where—under physiological and

pathological conditions, how purinergic transmitters are converted from one to another, and details regarding their spatial and temporal release patterns. Answering these questions requires a detection method that provides high spatial and temporal resolution, as well as high molecular precision in order to monitor the dynamics of structurally similar purinergic transmitters, ideally in living animals. Here, we review the technologies that are currently available for detecting purinergic transmitters, including classic non-optical methods and recently developed optical methods. The molecular, pharmacological, and functional profiles of purinergic signaling have been reviewed by other groups (Burnstock, 2008; Chen et al., 2013) and are not discussed here in detail.

2. Non-optical methods for probing purinergic transmitters

2.1. Electrophysiology

Because the binding of ATP to P2X receptors induces an increase in cationic conductance, electrophysiological recording can be used to measure ATP by recording currents through P2X receptors (Fig. 1A1). Using the patch-clamp technique, ATP-induced P2X currents can be recorded in cells expressing endogenous P2X receptors or in “sniffer cells” that express P2X receptors (Hayashi et al., 2004; Zhang et al., 2019). In addition to ionotropic receptors, all purinergic transmitters identified to date also have one or more corresponding metabotropic G protein-coupled receptors (GPCRs), which transduces receptor activation into downstream electrophysiological effects on target cells. For example, electrophysiological studies in guinea pig hippocampal slices revealed that NMDA-induced depression of evoked excitatory postsynaptic currents was abolished by the adenosine A₁ receptor antagonist 8-cyclo-pentyltheophylline, suggesting that adenosine is released in an activity-dependent manner (Manzoni et al., 1994). Similarly, the binding of non-adenosine purinergic transmitters to P2Y receptors induces the G protein-mediated opening or closing of downstream ion channels, giving rise to an electrophysiological change that can be recorded in target cells. Although electrophysiological methods are extremely sensitive, they typically have extremely low throughput. Moreover, because electrophysiological methods mainly detect the downstream effects of purinergic transmitters, pharmacology is often required in order to verify molecular specificity. Finally, although electrophysiology can achieve temporal resolution on the order of milliseconds, it does not provide a measure of the transmitter's release kinetics. Thus, combining electrophysiology with direct detection methods would enable researchers to study both the release of purinergic transmitters and their downstream effects in target cells.

2.2. Microdialysis

A widely used method for directly measuring purinergic transmitters is microdialysis (Fig. 1A2) (Bito et al., 1966; Porkka-Heiskanen et al., 1997), often in combination with high-performance liquid chromatography (HPLC) together with either ultraviolet (UV) absorption detector or mass spectrometry (MS). With this method, a microdialysis probe (~200 μm in diameter), containing physiological solution such as artificial cerebrospinal fluid (aCSF), is first implanted into the brain region(s) of interest (Porkka-Heiskanen et al., 2000), and extracellular small molecules such as purinergic transmitters are collected via passive diffusion through the semi-permeable membrane. The dialysates are then separated by HPLC and detected using MS or UV. Microdialysis has several clear advantages compared to other approaches. First, it can be used in living animals, including humans. Second, this approach has extremely high sensitivity for detecting purinergic transmitters




Fig. 1. Overview of select non-optical and optical methods for detecting purinergic transmitters.

In each panel, the principle (with example data, where indicated) behind for each method is shown at the left; the corresponding radar graph summarizing the method's performance index (including sensitivity, molecular specificity, cell-type, non-invasiveness, spatial resolution, and temporal resolution) is shown at the right, with arbitrary units ranging from 0 to 4. Abbreviations: aCSF, artificial cerebrospinal fluid; Ado, adenosine; ATP, adenosine triphosphate; CFP, cyan fluorescent protein; FRET, fluorescence resonance energy transfer; IgK, light chain kappa; MS, mass spectrometry; NMDAR, *N*-methyl-D-aspartate receptor; PDGFR, platelet-derived growth factor receptor; YFP, yellow fluorescent protein. Schematic shown in (A3) adapted from (Dale and Frenguelli, 2012); data shown in (A4) adapted from (Swamy and Venton, 2007); schematic shown in (B3) adapted from (Conley et al., 2017); schematic shown in (B4) adapted from (Lobas et al., 2019).

(in the nanomolar range). Third, it can be used to detect multiple chemicals simultaneously in one sample. Finally, it can provide the absolute concentrations of multiple chemicals. On the other hand, microdialysis also has several drawbacks. First, the sampling rate is relatively slow, on the order of minutes; this low temporal resolution precludes the possibility of capturing the rapid dynamics of purinergic transmitters. In addition, purinergic transmitters can be metabolized during the lengthy sample preparation. A second disadvantage of microdialysis is that although the microdialysis probe can be quite small, it cannot provide subcellular spatial resolution or even cell-type specificity. Finally, implanting microdialysis probe(s) in the brain can cause tissue damage and/or neuroinflammation, thereby potentially altering the concentration of purinergic transmitters in the local microenvironment (Ohyama et al., 2019).

2.3. Amperometric biosensors

In order to detect purinergic transmitters in real time, researchers have developed a variety of electrochemistry-based methods using electrodes, including enzyme-based amperometric biosensors (Dale, 1998; Dale et al., 2000) and fast-scan cyclic voltammetry (FSCV) using carbon-fiber microelectrodes (Swamy and Venton, 2007). The basic principle of amperometric biosensors (Fig. 1A3) capitalizes on the metabolism of purinergic transmitters to create hydrogen peroxide (H_2O_2) via an enzymatic cascade; the resulting H_2O_2 can then be detected using an amperometer at the microelectrode's surface (Dale and Frenguelli, 2012). For example, to detect ATP, the surface of the microelectrode is coated with the glycerol kinase and glycerol-3-phosphate oxidase (Laudet et al., 2005). In the presence of ATP, glycerol is phosphory-

lated by glycerol kinase to produce glycerol-3-phosphate, which is then oxidized by glycerol-3-phosphate oxidase, generating H_2O_2 . When glycerol is available at saturated levels, this biosensor provides a final readout that is proportional to the concentration of ATP. A similar strategy has been used to detect adenosine, in this case using three enzymes, adenosine deaminase, nucleotide phosphorylase, and xanthine oxidase (Dale and Frenguelli, 2012). During the enzymatic cascade, adenosine is first converted to inosine, then to hypoxanthine, and finally to xanthine, urate, and H_2O_2 .

Amperometric biosensors provide sufficient temporal resolution and nanomolar-range sensitivity for detecting ATP and/or adenosine dynamics in different brain regions, including the hippocampus (Wall and Dale, 2013) and cerebellum (Wall and Dale, 2007), and under a variety of physiological and pathological conditions (Dale and Frenguelli, 2009; Gourine et al., 2005). On the other hand, amperometric biosensors have relatively poor molecular specificity and can also detect intermediate products. For example, the adenosine amperometric biosensor is sensitive to both adenosine and its metabolite inosine. Therefore, a control experiment should be performed using a sensor that does not include the adenosine deaminase (Dale and Frenguelli, 2012). When measuring adenosine, the control sensor can be positioned close to the full biosensor in order to distinguish adenosine from similar compounds; the adenosine-specific signal can then be calculated by subtracting the control sensor's signal from the full biosensor's signal. Similarly, using an amperometric biosensor to measure ATP has this problem as well. In addition, as with other electrode-based sensors, placing the amperometric biosensors in the brain is invasive and can lead to tissue damage. Finally, amperometric biosensors usually have relatively limited spatial resolution and are unable to detect purinergic transmitter in specific cell types. In principle, if an enzymatic cascade catalyzes the bioconversion of purinergic transmitters other than adenosine or ATP, and if that cascade produces H_2O_2 at levels proportional to the original transmitter's concentration, it should be possible to develop new amperometric biosensors that can detect a diverse number of purinergic transmitters; however, to the best of our knowledge, this method has only been used to detect adenosine and ATP.

2.4. Fast-scan cyclic voltammetry

In addition to amperometry, fast-scan cyclic voltammetry (FSCV) with carbon-fiber microelectrodes (Fig. 1A4) has also been used as an electrochemical method to directly monitor purinergic transmitters, including adenosine (Swamy and Venton, 2007). Traditionally, FSCV has been used to detect monoamine neuromodulators such as dopamine, norepinephrine, and serotonin (Bunin et al., 1998; Park et al., 2011), as monoamines can be readily oxidized, thereby giving rise to signature currents under a specific cyclic voltage scan range. Because adenosine is also an electroactive molecule that undergoes a series of two-electron oxidation steps, it can also be detected using FSCV (Nguyen and Venton, 2015). Compared to amperometric biosensors, FSCV electrodes have similar sensitivity (i.e., in the nanomolar range). Importantly, however, FSCV can achieve temporal resolution on the order of hundreds of milliseconds; thus, FSCV has been used to detect rapid activity-dependent adenosine release (Pajski and Venton, 2010), as well as spontaneous transient adenosine signals (Nguyen et al., 2014) in brain slices. With respect to specificity, adenosine metabolites such as inosine are electrochemically inactive to carbon-fiber electrodes; therefore, FSCV has higher specificity than amperometric biosensors (Nguyen and Venton, 2015). Moreover, although FSCV was successfully used for monitoring adenosine, it was still difficult to expand this method for measuring other purinergic transmitters.

Finally, as with other electrode-based methods, FSCV lacks subcellular spatial resolution and cell-type specificity.

3. Optical methods

Compared to the conventional non-optical methods discussed above, optical imaging methods provide direct, sensitive, higher throughput, and long-term monitoring neurotransmitter dynamics in cultured cells, brain slices, and behaving animals *in vivo*. More importantly, the genetically-encoded optical sensor could be targeted to defined cell-types or subcellular compartments to achieve the cell-type specificity and/or subcellular spatial resolution. Compared with the classical electrophysiology recording, optical imaging methods are less invasive and therefore provide repeatable measurements of neurotransmitter dynamics from the same neuronal population. Therefore, optical methods are becoming increasingly popular. Several optical sensors have been developed for detecting both intracellular purine metabolites (Table 1) and extracellular purinergic transmitters (Table 2), and are discussed in further detail below.

3.1. Optical sensors for detecting intracellular purine metabolites

Within the cytoplasm, purine metabolites—particularly ATP—serve as a universal and highly important source of cellular energy, and several optical methods have been developed in order to investigate how ATP affects cellular physiology. For example, firefly luciferase—an ATP-consuming enzyme that oxidizes the small-molecule substrate luciferin to produce a bioluminescent product—has been widely used to measure ATP both *in vitro* and *in vivo*. Importantly, luciferase can be expressed in specific target cells or even in subcellular organelles when fused with sequences that target the protein to specific structures such as the mitochondria or synaptic vesicles (Rangaraju et al., 2014); organelle-specific luciferase-based sensors have been discussed in detail elsewhere (Morciano et al., 2017). Unfortunately, luciferase can only be used to detect ATP, not other purine metabolites. Moreover, by detecting ATP, the enzymatic reaction itself consumes ATP, which inevitably leads to changes in cellular metabolism. Recently, Mathew Tantama and colleagues developed a ratiometric bioluminescence resonance energy transfer (BRET) sensor for detecting ATP; this system, called ARSeNL, uses the ATP-independent NanoLuc luciferase and the red fluorescent protein mScarlet as the donor and acceptor, respectively (Min et al., 2019). Using this sensor, they measured changes in energy metabolism both *in vitro* and *in vivo*. Because mammalian cells do not normally express bioluminescent proteins, luminescence-based methods usually have an extremely low background signal (Morciano et al., 2017); on the other hand, bioluminescence-based methods produce low numbers of photons and lack cellular-scale resolution, which can limit its general applicability.

Unlike bioluminescence-based methods, sensors based on fluorescent proteins can—at least in principle—provide higher spatial and temporal resolution with respect to imaging purinergic transmitters. A fluorescent sensor requires two components: *i*) a binding protein that specifically recognizes purinergic transmitters and generates a conformational change upon ligand binding, and *ii*) a fluorescent component that converts the conformational change into a fluorescent signal. To develop such a sensor, several proteins were covalently coupled to a fluorescent reporter dye that undergoes a change in fluorescence upon binding ATP or ADP, thereby providing a measure of ATP/ADP concentration (Kunzelmann and Webb, 2009, 2010; Vancaenenbroeck and Webb, 2015). Such chemical dye-based methods provide an important tool for quantifying ATP with high sensitivity and high throughput; however,

Table 1
Overview of optical sensors for detecting intracellular purine metabolites.

Sensor	Ligand	Binding protein / module	Optical reporter	$\Delta F/F_0$ or FRET/BRET ratio	Kd or EC ₅₀	Kinetics	<i>In vivo</i> application	Refs.
Luciferase	ATP	No	Bioluminescence	N.D.	N.D.	N.D.	N.D.	DeLuca and McElroy (1974)
Syn-ATP	Presynaptic ATP	No	Bioluminescence	~0.25	2.3 mM	N.D.	N.D.	Rangaraju et al. (2014)
ParM-based sensor	ADP	Bacterial actin-like ParM	Chemical dyes	>3.5/~15	0.45/~30 μ M	$0.65/9.5 \times 10^4$ M ⁻¹ s ⁻¹ (k _{on}), 0.19/2.9 s ⁻¹ (k _{off})	N.D.	Kunzelmann and Webb (2009, 2010)
FRET-Based chemosensor	Nucleoside polyphosphates	Binuclear zinc complex	Chemical dyes	~33	0.1–1 μ M	N.D.	N.D.	Kurishita et al. (2010)
Perceval and PercevalHR	Cytosolic ATP/ADP ratio	Bacterial GlnK1	cpmVenus	K _R [*] : 0.5/3.5	N.D.	N.D.	N.D.	Berg et al. (2009); Tantama et al. (2013)
ATeam	Cytosolic ATP	Bacterial F ₀ F ₁ -ATP synthase	CFP/YFP	~1.5	7.4 μ M–3.3 mM	~17 μ M/s (τ_{on}), 98 ms (τ_{off})	N.D.	Imamura et al. (2009)
ARSeNL	ATP	Bacterial F ₀ F ₁ -ATP synthase	mScarlet/ NanoLuc	~5	~1.1 mM	N.D.	Mice	Min et al. (2019)
QUEEN	Cytosolic ATP	Bacterial F ₀ F ₁ -ATP synthase	cpEGFP	<4 (excitation ratio)	7 μ M	N.D.	N.D.	Yaginuma et al. (2014)
iATPSnFR	ATP	Bacterial F ₀ F ₁ -ATP synthase	cpSFGFP	~2.4/~1.9	~120 μ M/~50 μ M	N.D.	N.D.	Lobas et al. (2019)

K_R^{*} represents a half-maximal signal change, which was used to quantify the ability of Perceval sensors to report ATP/ADP ratio. N.D., not determined.

Table 2
Overview of optical sensors for detecting extracellular purinergic transmitters.

Sensor	Ligand	Binding protein / module	Optical reporter	$\Delta F/F_0$ or FRET/BRET ratio	Kd or EC ₅₀	Kinetics	<i>In vivo</i> application	Refs.
pmeLUC	ATP	No	Bioluminescence	~8	> 10 μ M	N.D.	Mice	Morciano et al. (2017)
TANGO Assay Sniffer cells	GPCR ligands Adenosine, ATP, or ADP	GPCRs A ₁ R, P2X, or P2Y	Reporter gene Ca ²⁺ sensor	>10 ~2.0 for Ado?	~ 1 nM 8 μ M for Ado	>8 h N.D.	N.D. N.D.	Barnea et al. (2008) Yamashiro et al. (2017)
ecATeam3.10	ATP	Bacterial F ₀ F ₁ -ATP synthase	CFP/YFP	~0.27	~11 mM	~13–172 s (τ_{on})	N.D.	Conley et al. (2017)
A _{2A} R-based FIAsh	Adenosine	A _{2A} R	CFP/FIAsh	~0.1	Similar to WT A _{2A} R ?	66–88 ms (τ_{on})	N.D.	Hoffmann et al. (2005)
P2X ₂ -Cam	ATP	P2X ₂	CFP/YFP	~0.37	~14 μ M	~6 s (decay)	Zebrafish	Richler et al. (2008)
iATPSnFR	ATP	Bacterial F ₀ F ₁ -ATP synthase	cpSFGFP	~1.0/~0.9	~350 μ M/ ~140 μ M	<5 s (τ_{on}), <4 s (τ_{off})	N.D.	Lobas et al. (2019)

N.D., not determined.

because it can be challenging to target these tools to specific cells, this method is generally limited.

To overcome this limitation, Gary Yellen and colleagues engineered Perceval—a fully genetically encoded fluorescent protein-based sensor for imaging ATP—by inserting circularly permuted monomeric Venus (cpmVenus) into GlnK1, an ATP-binding bacterial protein that regulates ammonia transport (Berg et al., 2009). Like the native GlnK1 protein, Perceval measures the ATP/ADP ratio. However, because Perceval binding is saturated at a relatively low ATP/ADP ratio (< 5:1), its effectiveness is limited in mammalian cells. They therefore increased the sensor's dynamic range so that it can still detect ATP/ADP ratios in healthy mammalian cells, which can reach 100:1. This optimized sensor, called PercevalHR (Tantama et al., 2013), produces a stronger signal, thus allowing researchers to measure the ATP/ADP ratio at the single-cell level; moreover, PercevalHR performs well using either one-photon or two-photon excitation. Unfortunately, however, both Perceval and PercevalHR are highly sensitive to pH; therefore, the fluorescence change that derives from the bona fide signal can be difficult to distinguish from an artifact. Moreover, the Perceval-based sensors can only be used to monitor cellular energy levels and would be difficult to use for directly quantifying specific purine metabolites.

To specifically visualize ATP levels within individual living cells, Hiroyuki Noji and colleagues generated a series of sensors based on fluorescence resonance energy transfer (FRET); these sensors, called ATeams, have specific apparent affinities ranging from 7.4 μ M to 3.3 mM (Imamura et al., 2009). ATeams contain the epsilon subunit of bacterial F₀F₁-ATP synthase as the ATP-sensing protein “sandwiched” between cyan fluorescent protein (CFP) and yellow fluorescent protein (YFP). By targeting these sensors to specific subcellular compartments, Noji's group successfully measured ATP levels within the mitochondrial matrix, cytoplasm, and nucleus in cultured HeLa cells. Unfortunately, because FRET efficiency is relatively low, the applications of ATeam are limited, particularly with *in vivo* preparations. In addition to their FRET-based ATeams, Hiroyuki Noji, Hiromi Imamura, and their colleagues developed an excitation ratio-based sensor called QUEEN by inserting circularly permuted enhanced green fluorescent protein (cpEGFP) between two α -helices of the bacterial F₀F₁-ATP synthase epsilon subunit; they then used this sensor to measure ATP levels in *E. coli* cells (Yaginuma et al., 2014). Despite their advantages, both the FRET-based ATeam sensors and the excitation ratio-based QUEEN sensors require customized equipment, have relatively slow imaging rates, and have a lower signal-to-noise ratio compared to single wavelength-based sensors (Lobas et al., 2019).

Recently, Baljit Khakh, Loren Looger, and their colleagues developed a genetically encoded GFP-based single-wavelength ATP sensor called iATPSnFR by inserting circularly permuted superfolder GFP (cpSFGFP) in the epsilon subunit of the *Bacillus PS3* F₀F₁-ATPase (Lobas et al., 2019). This newly developed sensor produces a bright, rapid, specific increase in fluorescence increase upon binding ATP (with peak $\Delta F/F_0$ values of ~3.9 using purified proteins); however, the low affinity of iATPSnFR for ATP (on the order of hundreds of micromolar) might not be sensitive enough for *in vivo* applications, particularly when measuring extracellular ATP.

3.2. Optical sensors for detecting extracellular purinergic transmitters

Several strategies have been used to develop optical sensors for detecting extracellular purinergic transmitters, including: *i*) targeting cytosolic sensors to the plasma membrane, and *ii*) using plasma membrane-localized proteins to engineer new optical sensors. The resulting optical sensors can be based on either bioluminescence or fluorescence, and are discussed in detail below.

3.2.1. Bioluminescence-based sensors

As discussed above, luciferase-based bioluminescent sensors are widely used for measuring ATP (Fig. 1B1). To measure local ATP release from activated platelets, George Dubyak and colleagues attached a purified IgG binding domain-tagged luciferase to the surface of cells expressing IgG (Beigi et al., 1999). Similarly, using luciferase either diluted in extracellular solution (Wang et al., 2000) or immobilized to beads (Zhang et al., 2008), Edward Yeung and colleagues successfully measured mechanical stimulation-evoked release of ATP from astrocytes *in vitro*. In addition, by fusing a leader sequence and glycosylphosphatidylinositol (GPI) lipid anchor derived from the folate receptor to luciferase, Francesco Di Virgilio, Paolo Pinton, and their colleagues designed a plasma membrane-targeted luciferase called pmeLUC, which they then used to measure extracellular ATP dynamics both in cultured cells and in living animals injected with pmeLUC-expressing cells (Morciano et al., 2017). Again, although luciferase-based bioluminescent sensors have an extremely low background signal, their output yields relatively low numbers of photons, which reduces their feasibility for imaging purinergic transmitters at the cellular scale. Interestingly, the bioluminescent signal can be amplified by attaching the sensor to a downstream GPCR (Barnea et al., 2008); however, these so-called TANGO assays usually have poor temporal resolution (on the order of minutes to hours), making them less feasible for monitoring purinergic transmitters in real time.




Fig. 2. A proposed toolbox for detecting purinergic transmitters.

(A) The principle behind our proposed GPCR Activation-Based (GRAB) sensors (left) for detecting transmitters, with the corresponding theoretical performance index (right). (B) A similar GRAB strategy was successfully used for *in vivo* monitoring of acetylcholine (ACh), dopamine (DA), and norepinephrine (NE) in behaving animals. (C) A proposed toolbox of GRAB sensors for monitoring various purinergic transmitters; the structure of each transmitter is shown at the top. Data shown in (B) adapted from (Jing et al., 2018), (Sun et al., 2018), (Patriarchi et al., 2018), and (Feng et al., 2019).

3.2.2. Fluorescence-based sensors

Compared to bioluminescence-based sensors, fluorescence-based sensors have significantly higher brightness, which enables researchers to visualize purinergic transmitters with optical precision and/or millisecond temporal resolution. Therefore, fluorescence-based sensors are often used to analyze molecules and events that occur with rapid dynamics at a subcellular level using both *in vitro* and *in vivo* preparations. Here, we discuss the recent progress in fluorescence-based sensors for measuring purinergic transmitters, including cell-based sensors, FRET-based sensors, and single-wavelength sensors.

3.2.2.1. Cell-based sensors.

Several cell-based sensors—also referred to as “sniffer” cells (Fig. 1B2)—have been developed for monitoring purinergic transmitters. With this method, a specific cell surface receptor that recognizes the purinergic transmitter of interest is expressed in the sniffer cells, which are then placed in close proximity to the target cells. For example, the Ca^{2+} dye Fluo-4-AM was loaded in the mouse glioma cell line GL261, which express high-sensitivity purinergic receptors, and these cells were then placed on the surface of a mouse brain slice and used to indirectly image ATP release; the Ca^{2+} signal measured in the sniffer cells was due specifically to the release of ATP, as it was

blocked by the purinergic receptor blocker Reactive Blue-2 (Haas et al., 2006). A similar strategy was used to detect changes in extracellular adenosine by placing A₁R-expressing and Fura-2 loaded HEK293 T cells in the brain slice (Yamashiro et al., 2017). Although cell-based sensors can report specific purinergic transmitters with high sensitivity, they are not suitable for detecting the release of purinergic transmitters with high cell-type specificity or subcellular resolution. In addition, the sniffer cells would need to be implanted in specific tissues when used in living animals, making this approach potentially too invasive for use in *in vivo* applications.

3.2.2.2. FRET-based sensors. In principle, the above-mentioned limitations can be overcome using a genetically encoded probe, which allows researchers to monitor neuromodulator dynamics in specific cell types with subcellular spatial resolution. One such example is FRET-based sensors. For example, Martin Lohse and colleagues developed a CFP/FlAsH-based FRET sensor using the adenosine 2A receptor (A_{2A}R) as the backbone (Hoffmann et al., 2005). This sensor, which has similar affinity for A_{2A}R agonists compared to the wild-type A_{2A}R, produces an ~10 % change in the FRET ratio in response to a saturated concentration of adenosine. Similarly, Baljit Khakh and colleagues used the ATP receptor P2X₂ as the detecting module in their FRET sensor P2X₂-Cam for imaging ATP; specifically, they fused the P2X₂ receptor with a YC3.1 tag, which contains CFP and YFP separated by the Ca²⁺ sensor Cameleon (Richler et al., 2008). The Cameleon domain undergoes a conformational change upon binding Ca²⁺, changing the relative positions of the CFP and YFP moieties, thereby causing an increase in the FRET ratio. The authors found that P2X₂-Cam has an ~37 % change in the FRET ratio in response to a saturated concentration of ATP, with an EC₅₀ of approximately 13 μM. Unfortunately, because FRET-based sensors have a relatively low signal-to-noise ratio, they are less suitable for use in *in vivo* applications. Indeed, although P2X₂-Cam was used in live zebrafish to measure the application of ATP-induced ratio-metric changes, it has not been used to detect endogenous ATP release *in vivo*. In addition to using plasma membrane-localized A_{2A}R or P2X₂ receptors to engineer optical sensors, cytosolic sensors could also be targeted to the plasma membrane for probing extracellular purinergic transmitters. For example, Mathew Tantama and colleagues re-engineered the FRET-based ATeam ATP sensor to be expressed on the cell surface, and the new ecAT3.10 sensor (Fig. 1B3) enabled probing ATP release in cultured cells but not *in vivo* (Conley et al., 2017).

3.2.2.3. Single-wavelength sensors. Compared to FRET-based sensors, single-wavelength sensors have a higher signal-to-noise ratio and are therefore more suitable for measuring endogenous neuromodulator dynamics *in vivo*. For example, in order to image extracellular ATP, the genetically encoded single-wavelength iATP-SnFR (Fig. 1B4) was displayed at the extracellular side of the plasma membrane (Lobas et al., 2019). Despite their advantages, however, sensors based on bacterial binding proteins—including iATPSnFR—have relatively low ligand affinity, which limits their application, particularly with respect to *in vivo* preparations. Although the sensor's affinity could theoretically be improved through molecular engineering, this can be highly time-consuming. Moreover, although this strategy has been used to develop ATP sensors, to the best of our knowledge it has not been used to develop sensors for other purinergic transmitters.

4. Future perspectives

Despite significant advances over the past few decades, current methods for detecting purinergic transmitters are still limited in

several respects, including poor spatial and/or temporal resolution, insufficient sensitivity, and/or low specificity; thus, a method that satisfies all of these criteria is needed. Yet the question remains: how to engineer a sensor that can be used to monitor structurally similar purinergic transmitters with high specificity and sensitivity? In fact, nature has already provided us with an important clue: in mammals, all purinergic transmitters identified to date bind at least one G protein-coupled receptor. Thus, GPCRs provide an excellent scaffold for developing tools to detect purinergic transmitters. In this respect, a promising new strategy is to generate single-fluorophore GPCR Activation-Based (GRAB) sensors (Jing et al., 2019); indeed, this strategy has been used successfully to engineer sensors for detecting acetylcholine (Fig. 2A) (Jing et al., 2018), dopamine (Patriarchi et al., 2018; Sun et al., 2018), and norepinephrine (Feng et al., 2019). Importantly, these sensors have a wide dynamic range, rapid kinetics, single-cell resolution, and—most importantly—exquisite selectivity (Fig. 2B). In addition, most GRAB sensors have negligible coupling with their GPCR's downstream signaling pathways, and have no detectable effects on cellular physiology (Wu et al., 2019). These sensors allow researchers to monitor neurotransmitter dynamics *in vivo* in a variety of organisms, including flies, fish, birds (Tanaka et al., 2018), and mice (Fig. 2C).

Given the wide variety of chemical-sensing GPCRs expressed in the nervous system, this strategy can likely be used to develop sensors for detecting a wide range of molecules, including purinergic transmitters (Fig. 2D). Moreover, by attaching fluorescent proteins with different excitation/emission profiles, multiple sensors with non-overlapping spectra can be developed and then used in combination to study the complex interplay between different purinergic transmitters. In this respect, the use of simultaneous multi-color imaging to measure the release of multiple purinergic transmitters in real time will provide important insights into how these purinergic transmitters function in the brain and other tissues in both health and disease.

Author contributions

Y. L. and Z. W. conceived and wrote the manuscript.

Acknowledgments and Declaration of Competing Interest

We thank the members of the Li lab for providing feedback on the manuscript. This work was supported by the Beijing Municipal Science & Technology Commission (Z181100001318002), the Beijing Brain Initiative of Beijing Municipal Science & Technology Commission (Z181100001518004), the Guangdong grant 'Key technologies for treatment of brain disorders' (grant 2018B030332001), the General Program of National Natural Science Foundation of China (project 31671118 and project 31871074), the NIH BRAIN Initiative (NS103558), grants from the Peking-Tsinghua Center for Life Sciences and the State Key Laboratory of Membrane Biology at Peking University School of Life Sciences (to Y. L.). Z. W. is supported by the Boehringer Ingelheim-Peking University Postdoctoral Fellowship. Y. L. has filed patent applications, the value of which may be affected by this publication.

References

- Barnea, G., Strapps, W., Herrada, G., Berman, Y., Ong, J., Kloss, B., Axel, R., Lee, K.J., 2008. The genetic design of signaling cascades to record receptor activation. *Proc. Natl. Acad. Sci. U. S. A.* 105, 64–69.
- Beigi, R., Kobatake, E., Aizawa, M., Dubyak, G.R., 1999. Detection of local ATP release from activated platelets using cell surface-attached firefly luciferase. *Am. J. Physiol.-Cell Physiol.* 276, C267–C278.
- Berg, J., Hung, Y.P., Yellen, G., 2009. A genetically encoded fluorescent reporter of ATP:ADP ratio. *Nat. Methods* 6, 161–166.

- Bito, L., Davson, H., Levin, E., Murray, M., Snider, N., 1966. The concentrations of free amino acids and other electrolytes in cerebrospinal fluid, in vivo dialysate of brain, and blood plasma of the dog. *J. Neurochem.* 13, 1057–1067.
- Boison, D., 2012. Adenosine dysfunction in epilepsy. *Glia* 60, 1234–1243.
- Bunin, M.A., Prioleau, C., Mailman, R.B., Wightman, R.M., 1998. Release and uptake rates of 5-hydroxytryptamine in the dorsal raphe and substantia nigra reticulata of the rat brain. *J. Neurochem.* 70, 1077–1087.
- Burnstock, G., 1972. Purinergic nerves. *Pharmacol. Rev.* 24, 509–581.
- Burnstock, G., 2007. Physiology and pathophysiology of purinergic neurotransmission. *Physiol. Rev.* 87, 659–797.
- Burnstock, G., 2008. Purinergic signalling and disorders of the central nervous system. *Nat. Rev. Drug Discov.* 7, 575.
- Chen, J.-F., Eltzschig, H.K., Fredholm, B.B., 2013. Adenosine receptors as drug targets—what are the challenges? *Nat. Rev. Drug Discov.* 12, 265.
- Conley, J.M., Radhakrishnan, S., Valentino, S.A., Tantama, M., 2017. Imaging extracellular ATP with a genetically-encoded, ratiometric fluorescent sensor. *PLoS One* 12, e0187481.
- Corti, F., Cellai, L., Melani, A., Donati, C., Bruni, P., Pedata, F., 2013. Adenosine is present in rat brain synaptic vesicles. *Neuroreport* 24, 982–987.
- Cotrina, M.L., Lin, J.H., Alves-Rodrigues, A., Liu, S., Li, J., Azmi-Ghadimi, H., Kang, J., Naus, C.C., Nedergaard, M., 1998. Connexins regulate calcium signaling by controlling ATP release. *Proc. Natl. Acad. Sci. U. S. A.* 95, 15735–15740.
- Dale, N., 1998. Delayed production of adenosine underlies temporal modulation of swimming in frog embryo. *J. Physiol.* 511 (Pt 1), 265–272.
- Dale, N., Frenguelli, B.G., 2009. Release of adenosine and ATP during ischemia and epilepsy. *Curr. Neuropharmacol.* 7, 160–179.
- Dale, N., Frenguelli, B.G., 2012. Measurement of purine release with microelectrode biosensors. *Purinergic Signal.* 8, 27–40.
- Dale, N., Pearson, T., Frenguelli, B.G., 2000. Direct measurement of adenosine release during hypoxia in the CA1 region of the rat hippocampal slice. *J. Physiol.* 526 (Pt 1), 143–155.
- DeLuca, M., McElroy, W.D., 1974. Kinetics of the firefly luciferase catalyzed reactions. *Biochemistry* 13, 921–925.
- Dias, R.B., Rombo, D.M., Ribeiro, J.A., Henley, J.M., Sebastiao, A.M., 2013. Adenosine: setting the stage for plasticity. *Trends Neurosci.* 36, 248–257.
- Drury, A., Szent-Györgyi, A., 1929. The physiological activity of adenine compounds with especial reference to their action upon the mammalian heart 1. *J. Physiol.* 68, 213–237.
- Dunwiddie, T.V., 1999. Adenosine and Ethanol, The “Drunken” Synapse. Springer, pp. 119–133.
- During, M.J., Spencer, D.D., 1992. Adenosine: a potential mediator of seizure arrest and postictal refractoriness. *Ann. Neurol.* 32, 618–624.
- Eltzschig, H.K., Eckle, T., 2011. Ischemia and reperfusion—from mechanism to translation. *Nat. Med.* 17, 1391.
- Eltzschig, H.K., Sitkovsky, M.V., Robson, S.C., 2012. Purinergic signaling during inflammation. *N. Engl. J. Med.* 367, 2322–2333.
- Engel, T., Alves, M., Sheedy, C., Henshall, D.C., 2016. ATPergic signalling during seizures and epilepsy. *Neuropharmacology* 104, 140–153.
- Feng, J., Zhang, C., Lischinsky, J.E., Jing, M., Zhou, J., Wang, H., Zhang, Y., Dong, A., Wu, Z., Wu, H., Chen, W., Zhang, P., Zou, J., Hires, S.A., Zhu, J.J., Cui, G., Lin, D., Du, J., Li, Y., 2019. A genetically encoded fluorescent sensor for rapid and specific in vivo detection of norepinephrine. *Neuron* 102, 745–761, e748.
- Gourine, A.V., Llaudet, E., Dale, N., Spyer, K.M., 2005. ATP is a mediator of chemosensory transduction in the central nervous system. *Nature* 436, 108.
- Haas, B., Schipke, C.G., Peters, O., Sohl, G., Willecke, K., Kettenmann, H., 2006. Activity-dependent ATP-waves in the mouse neocortex are independent from astrocytic calcium waves. *Cereb. Cortex* 16, 237–246.
- Hasko, G., Linden, J., Cronstein, B., Pacher, P., 2008. Adenosine receptors: therapeutic aspects for inflammatory and immune diseases. *Nat. Rev. Drug Discov.* 7, 759–770.
- Hayashi, S., Hazama, A., Dutta, A.K., Sabirov, R.Z., Okada, Y., 2004. Detecting ATP release by a biosensor method. *Sci. STKE* 2004, pl14.
- Hockley, J.R., Tranter, M.M., McGuire, C., Boundouki, G., Cibert-Goton, V., Thaha, M.A., Blackshaw, L.A., Michael, J.E., Baker, M.D., Knowles, C.H., 2016. P2Y receptors sensitize mouse and human colonic nociceptors. *J. Neurosci.* 36, 2364–2376.
- Hoffmann, C., Gaietta, G., Bunemann, M., Adams, S.R., Oberdorff-Maass, S., Behr, B., Vilardaga, J.P., Tsien, R.Y., Ellisman, M.H., Lohse, M.J., 2005. A FAsH-based FRET approach to determine G protein-coupled receptor activation in living cells. *Nat. Methods* 2, 171–176.
- Holton, F.A., Holton, P., 1954. The capillary dilator substances in dry powders of spinal roots: a possible role of adenosine triphosphate in chemical transmission from nerve endings. *J. Physiol.* 126, 124–140.
- Holton, P., 1959. The liberation of adenosine triphosphate on antidromic stimulation of sensory nerves. *J. Physiol.* 145, 494–504.
- Huang, Z.-L., Qu, W.-M., Eguchi, N., Chen, J.-F., Schwarzschild, M.A., Fredholm, B.B., Urade, Y., Hayashi, O., 2005. Adenosine A_{2A}, but not A₁, receptors mediate the arousal effect of caffeine. *Nat. Neurosci.* 8, 858.
- Imamura, H., Nhat, K.P., Togawa, H., Saito, K., Iino, R., Kato-Yamada, Y., Nagai, T., Noji, H., 2009. Visualization of ATP levels inside single living cells with fluorescence resonance energy transfer-based genetically encoded indicators. *Proc. Natl. Acad. Sci. U. S. A.* 106, 15651–15656.
- Jin, J., Daniel, J.L., Kunapuli, S.P., 1998. Molecular basis for ADP-induced platelet activation II. The P2Y₁ receptor mediates ADP-induced intracellular calcium mobilization and shape change in platelets. *J. Biol. Chem.* 273, 2030–2034.
- Jing, M., Zhang, P., Wang, G., Feng, J., Mesik, L., Zeng, J., Jiang, H., Wang, S., Looby, J.C., Guagliardo, N.A., Langma, L.W., Lu, J., Zuo, Y., Talmage, D.A., Role, L.W., Barrett, P.Q., Zhang, L.L., Luo, M., Song, Y., Zhu, J.J., Li, Y., 2018. A genetically encoded fluorescent acetylcholine indicator for in vitro and in vivo studies. *Nat. Biotechnol.* 36, 726–737.
- Jing, M., Zhang, Y., Wang, H., Li, Y., 2019. G-protein-coupled receptor-based sensors for imaging neurochemicals with high sensitivity and specificity. *J. Neurochem.* 151, 279–288.
- Jo, Y.H., Schlichter, R., 1999. Synaptic corelease of ATP and GABA in cultured spinal neurons. *Nat. Neurosci.* 2, 241–245.
- Khakh, B.S., North, R.A., 2006. P2X receptors as cell-surface ATP sensors in health and disease. *Nature* 442, 527.
- Kunzelmann, S., Webb, M.R., 2009. A biosensor for fluorescent determination of ADP with high time resolution. *J. Biol. Chem.* 284, 33130–33138.
- Kunzelmann, S., Webb, M.R., 2010. A fluorescent, reagentless biosensor for ADP based on tetramethylrhodamine-labeled ParM. *ACS Chem. Biol.* 5, 415–425.
- Kurishita, Y., Kohira, T., Ojida, A., Hamachi, I., 2010. Rational design of FRET-based ratiometric chemosensors for in vitro and in cell fluorescence analyses of nucleoside polyphosphates. *J. Am. Chem. Soc.* 132, 13290–13299.
- Lazarowski, E.R., Harden, T.K., 2015. UDP-sugars as extracellular signaling molecules: cellular and physiologic consequences of P2Y₁₄ receptor activation. *Mol. Pharmacol.* 88, 151–160.
- Li, N., Lu, Z., Yu, L., Burnstock, G., Deng, X., Ma, B., 2014. Inhibition of G protein-coupled P2Y₂ receptor induced analgesia in a rat model of trigeminal neuropathic pain. *Mol. Pain* 10, 21.
- Llaudet, E., Hatz, S., Droniou, M., Dale, N., 2005. Microelectrode biosensor for real-time measurement of ATP in biological tissue. *Anal. Chem.* 77, 3267–3273.
- Lobas, M.A., Tao, R., Nagai, J., Kronschlager, M.T., Borden, P.M., Marvin, J.S., Looger, L.L., Khakh, B.S., 2019. A genetically encoded single-wavelength sensor for imaging cytosolic and cell surface ATP. *Nat. Commun.* 10, 711.
- Lovatt, D., Xu, Q., Liu, W., Takano, T., Smith, N.A., Schnermann, J., Tieu, K., Nedergaard, M., 2012. Neuronal adenosine release, and not astrocytic ATP release, mediates feedback inhibition of excitatory activity. *Proc. Natl. Acad. Sci. U. S. A.* 109, 6265–6270.
- Ma, Z., Taruno, A., Ohmoto, M., Jyotaki, M., Lim, J.C., Miyazaki, H., Niisato, N., Marunaka, Y., Lee, R.J., Hoff, H., 2018. CALHM3 is essential for rapid ion channel-mediated purinergic neurotransmission of GPCR-mediated tastes. *Neuron* 98, 547–561, e510.
- MacDonald, P.E., Braun, M., Galvanovskis, J., Rorsman, P., 2006. Release of small transmitters through kiss-and-run fusion pores in rat pancreatic β cells. *Cell Metab.* 4, 283–290.
- Maienschein, V., Marxen, M., Volkmandt, W., Zimmermann, H., 1999. A plethora of presynaptic proteins associated with ATP-storing organelles in cultured astrocytes. *Glia* 26, 233–244.
- Manzoni, O.J., Manabe, T., Nicoll, R.A., 1994. Release of adenosine by activation of NMDA receptors in the hippocampus. *Science* 265, 2098–2101.
- Min, S.H., French, A.R., Trull, K.J., Tat, K., Varney, S.A., Tantama, M., 2019. Ratiometric BRET measurements of ATP with a genetically-encoded luminescent sensor. *Sensors (Basel)* 19, 3502.
- Morciano, G., Sarti, A.C., Marchi, S., Missiroli, S., Falzoni, S., Raffaghello, L., Pistoia, V., Giorgi, C., Di Virgilio, F., Pinton, P., 2017. Use of luciferase probes to measure ATP in living cells and animals. *Nat. Protoc.* 12, 1542.
- Nguyen, M.D., Lee, S.T., Ross, A.E., Ryals, M., Choudhry, V.I., Venton, B.J., 2014. Characterization of spontaneous, transient adenosine release in the caudate-putamen and prefrontal cortex. *PLoS One* 9, e87165.
- Nguyen, M.D., Venton, B.J., 2015. Fast-scan cyclic voltammetry for the characterization of rapid adenosine release. *Comput. Struct. Biotechnol. J.* 13, 47–54.
- Ohyama, K., Ikeshita, Y., Fuchigami, Y., Kawakami, S., Nakashima, M.N., Nakashima, M., 2019. Proteomic analysis and ATP assay reveal a positive effect of artificial cerebral spinal fluid perfusion following microdialysis sampling on repair of probe-induced brain damage. *J. Neurosci. Methods* 315, 1–5.
- Pajski, M.L., Venton, B.J., 2010. Adenosine release evoked by short electrical stimulations in striatal brain slices is primarily activity dependent. *ACS Chem. Neurosci.* 1, 775–787.
- Pankratov, Y., Lalo, U., Verkhratsky, A., North, R.A., 2006. Vesicular release of ATP at central synapses. *PLoS. Arch.* 452, 589–597.
- Park, J., Takmakov, P., Wightman, R.M., 2011. In vivo comparison of norepinephrine and dopamine release in rat brain by simultaneous measurements with fast-scan cyclic voltammetry. *J. Neurochem.* 119, 932–944.
- Patriarchi, T., Cho, J.R., Merten, K., Howe, M.W., Marley, A., Xiong, W.H., Folk, R.W., Broussard, G.J., Liang, R., Jang, M.J., Zhong, H., Dombek, D., von Zastrow, M., Nimmerjahn, A., Gradinaru, V., Williams, J.T., Tian, L., 2018. Ultrafast neuronal imaging of dopamine dynamics with designed genetically encoded sensors. *Science* 360, eaat4422.
- Porkka-Heiskanen, T., Strecker, R.E., McCarley, R.W., 2000. Brain site-specificity of extracellular adenosine concentration changes during sleep deprivation and spontaneous sleep: an in vivo microdialysis study. *Neuroscience* 99, 507–517.
- Porkka-Heiskanen, T., Strecker, R.E., Thakkar, M., Bjorkum, A.A., Greene, R.W., McCarley, R.W., 1997. Adenosine: a mediator of the sleep-inducing effects of prolonged wakefulness. *Science* 276, 1265–1268.
- Ralevic, V., Burnstock, G., 1998. Receptors for purines and pyrimidines. *Pharmacol. Rev.* 50, 413–492.
- Rangaraju, V., Calloway, N., Ryan, T.A., 2014. Activity-driven local ATP synthesis is required for synaptic function. *Cell* 156, 825–835.
- Ren, C., Gan, X., Wu, J., Qiu, C.-Y., Hu, W.-P., 2016. Enhancement of acid-sensing ion channel activity by metabotropic P2Y₁ receptors in primary sensory neurons. *Purinergic Signal.* 12, 69–78.

- Richler, E., Chaumont, S., Shigetomi, E., Sagasti, A., Khakh, B.S., 2008. Tracking transmitter-gated P2X cation channel activation in vitro and in vivo. *Nat. Methods* 5, 87–93.
- Sabirov, R.Z., Okada, Y., 2005. ATP release via anion channels. *Purinergic Signal.* 1, 311–328.
- Schwarzschild, M.A., Agnati, L., Fuxe, K., Chen, J.F., Morelli, M., 2006. Targeting adenosine A2A receptors in Parkinson's disease. *Trends Neurosci.* 29, 647–654.
- Shryock, J.C., Belardinelli, L., 1997. Adenosine and adenosine receptors in the cardiovascular system: biochemistry, physiology, and pharmacology. *Am. J. Cardiol.* 79, 2–10.
- Sun, F., Zeng, J., Jing, M., Zhou, J., Feng, J., Owen, S.F., Luo, Y., Li, F., Wang, H., Yamaguchi, T., Yong, Z., Gao, Y., Peng, W., Wang, L., Zhang, S., Du, J., Lin, D., Xu, M., Kreitzer, A.C., Cui, G., Li, Y., 2018. A genetically encoded fluorescent sensor enables rapid and specific detection of dopamine in flies, fish, and mice. *Cell* 174, 481–496, e419.
- Swamy, B.K., Venton, B.J., 2007. Subsecond detection of physiological adenosine concentrations using fast-scan cyclic voltammetry. *Anal. Chem.* 79, 744–750.
- Tanaka, M., Sun, F., Li, Y., Mooney, R., 2018. A mesocortical dopamine circuit enables the cultural transmission of vocal behaviour. *Nature* 563, 117.
- Tantama, M., Martinez-Francois, J.R., Mongeon, R., Yellen, G., 2013. Imaging energy status in live cells with a fluorescent biosensor of the intracellular ATP-to-ADP ratio. *Nat. Commun.* 4, 2550.
- Taruno, A., Vingtdeux, V., Ohmoto, M., Ma, Z., Dvoryanchikov, G., Li, A., Adrien, L., Zhao, H., Leung, S., Abernethy, M., 2013. CALHM1 ion channel mediates purinergic neurotransmission of sweet, bitter and umami tastes. *Nature* 495, 223.
- Vancraenenbroeck, R., Webb, M.R., 2015. A fluorescent, reagentless biosensor for ATP, based on malonyl-coenzyme a synthetase. *ACS Chem. Biol.* 10, 2650–2657.
- Wall, M.J., Dale, N., 2007. Auto-inhibition of rat parallel fibre-Purkinje cell synapses by activity-dependent adenosine release. *J. Physiol.* 581, 553–565.
- Wall, M.J., Dale, N., 2013. Neuronal transporter and astrocytic ATP exocytosis underlie activity-dependent adenosine release in the hippocampus. *J. Physiol.* 591, 3853–3871.
- Wang, Z., Haydon, P.G., Yeung, E.S., 2000. Direct observation of calcium-independent intercellular ATP signaling in astrocytes. *Anal. Chem.* 72, 2001–2007.
- Wu, Z., Feng, J., Jing, M., Li, Y., 2019. G protein-assisted optimization of GPCR-activation based (GRAB) sensors. *Neural Imaging and Sensing 2019.* 10865 Vol. International Society for Optics and Photonics, 108650N.
- Yaginuma, H., Kawai, S., Tabata, K.V., Tomiyama, K., Kakizuka, A., Komatsuzaki, T., Noji, H., Imamura, H., 2014. Diversity in ATP concentrations in a single bacterial cell population revealed by quantitative single-cell imaging. *Sci. Rep.* 4, 6522.
- Yamashiro, K., Fujii, Y., Maekawa, S., Morita, M., 2017. Multiple pathways for elevating extracellular adenosine in the rat hippocampal CA 1 region characterized by adenosine sensor cells. *J. Neurochem.* 140, 24–36.
- Zhang, J.M., Wang, H.K., Ye, C.Q., Ge, W., Chen, Y., Jiang, Z.L., Wu, C.P., Poo, M.M., Duan, S., 2003. ATP released by astrocytes mediates glutamatergic activity-dependent heterosynaptic suppression. *Neuron* 40, 971–982.
- Zhang, Q., Liu, B., Wu, Q., Liu, B., Li, Y., Sun, S., Wang, Y., Wu, X., Chai, Z., Jiang, X., Liu, X., Hu, M., Wang, Y., Yang, Y., Wang, L., Kang, X., Xiong, Y., Zhou, Y., Chen, X., Zheng, L., Zhang, B., Wang, C., Zhu, F., Zhou, Z., 2019. Differential co-release of two neurotransmitters from a vesicle fusion pore in mammalian adrenal chromaffin cells. *Neuron* 102, 173–183, e174.
- Zhang, Y., Phillips, G.J., Li, Q., Yeung, E.S., 2008. Imaging localized astrocyte ATP release with firefly luciferase beads attached to the cell surface. *Anal. Chem.* 80, 9316–9325.
- Zhang, Z., Chen, G., Zhou, W., Song, A., Xu, T., Luo, Q., Wang, W., Gu, X., Duan, S., 2007. Regulated ATP release from astrocytes through lysosome exocytosis. *Nat. Chem. Biol.* 9, 945.
- Zimmermann, H., Zebisch, M., Strater, N., 2012. Cellular function and molecular structure of ecto-nucleotidases. *Purinergic Signal.* 8, 437–502.